USING NORMALIZING AND OPT-OUT LANGUAGE



Syphilis Screening in Clients at Increased Risk

To keep you healthy, we recommend screening for syphilis, which is a common infection that may not have symptoms. To make sure we use the correct test, can you tell me if you have ever had syphilis in the past?



— If Client Answers NO

If Client is UNSURE

Ask for the location, date/year, and medication received. Do not perform POC testing—order lab-based syphilis test.

Perform POC Testing

Ask if they ever received two penicillin injections in the hip. If they cannot remember, perform POC testing, and if reactive, send confirmation to lab.

CTC-SRH.ORG: Training the Nation's Title X Workforce

This project is supported by 1 FPTPA006031-03-00 issued by the Office of the Assistant Secretary for Health of the U.S. Department of Health and Human Services (HHS) as part of a financial assistance award totaling \$1,000,000 with 100 percent funded by the Office of Population Affairs/OASH/HHS.

USING NORMALIZING AND OPT-OUT LANGUAGE



Syphilis Screening in Communities with High Prevalence

We recommend a test for syphilis to everyone because of the high rates in our community. To make sure we use the correct test, can you tell me if you have ever had syphilis in the past?





If Client is UNSURE

Ask for the location, date/year, and medication received. Do not perform POC testing—order lab-based syphilis test.

Perform POC Testing

Ask if they ever received two penicillin injections in the hip. If they cannot remember, perform POC testing, and if reactive, send confirmation to lab.

CTC-SRH.ORG: Training the Nation's Title X Workforce

This project is supported by 1 FPTPA006031-03-00 issued by the Office of the Assistant Secretary for Health of the U.S. Department of Health and Human Services (HHS) as part of a financial assistance award totaling \$1,000,000 with 100 percent funded by the Office of Population Affairs/OASH/HHS.